

VapoMeter

Transepidermal water loss TEWL, Skin barrier function, Penetration studies

A Novel Chimeric Lysin Shows Superiority to Mupirocin for Skin Decolonization of Methicillin-Resistant and -Sensitive *Staphylococcus aureus* Strains.
Pastagia M., Euler C., Chahales P., Fuentes-Ducalan J., Krueger J.G. and Fischetti V.A.
Antimicrob Agents Chemother. Feb; 55(2): 738-744, 2011.

Quantification of activated and total caspase-14 with newly developed ELISA systems in normal and atopic skin.
Yanamoto M., Kamata Y., Iida T., Fukushima H., Nomura J., Saito M., Tajima M., Okubo Y., Momoi T., Tsuboi R. and Hibino T.
J Dermatol Sci. Feb; 61(2): 110-117, 2011.

Documentation of normal stratum corneum scaling in an average population: features of differences among age, ethnicity and body site.
Chu M. and Kollias N.
Br J Dermatol. published online Jan 28 2011.

Formation and Closure of Microchannels in Skin Following Microporation.
Kalluri H. and Banga A.K.
Pharm Res. Jan; 28(1): 82-94, 2011.

Epidermal hyperproliferation in mice lacking fatty acid transport protein 4 (FATP4) involves ectopic EGF receptor and STAT3 signaling.
Lin M.-H., Chang K.-W., Lin S.-C. and Miner J.H.
Developmental Biology, Aug 15; 344(2): 707-719, 2010.

Iontophoresis mediated in vivo intradermal delivery of terbinafine hydrochloride.
Sancheva V., Kim H.D., Friden P.M. and Banga A.K.
Int J Pharm. Jun 30; 393(1-2), 113-119, 2010.

Comparison of tape stripping with the human skin blanching assay for the bioequivalence assessment of topical clobetasol propionate formulations.
Au W.L., Skinner M.F. and Kander I.
J Pharm Pharmaceut Sci. 13(1), 11-20, 2010.

The impact of obesity on skin disease and epidermal permeability barrier status.
Guida B., Nino M., Perrino N., Laccetti R., Trio R., Labella S. and Balato N.
J Eur Acad Dermatol Venereol. Feb; 24(2): 191-195, 2010.

Up-regulation of serpin SCCA1 is associated with epidermal barrier disruption.
Katagiri C., Iida T., Nakanishi J., Ozawa M., Aiba S. and Hibino T.
J Dermatol Sci. Feb; 57(2): 95-101, 2010.

Effects of Alcohol Intake on the Skin Physiology.
Jang W.S., Kim C.W., Kim S.E., Kim B.J. and Kim M.N.
Korean J Dermatol. Nov; 48(11): 948-954, 2010. Korean.

Tight junctions in Hailey-Hailey and Darier's diseases.
Raiko L., Leinonen P., Hägg P.M., Peltonen J., Oikarinen A. and Peltonen S.
Dermatology Reports, 1:e1, 2009

Functional Characteristics of the Skin Surface of Children Approaching Puberty: Age and Seasonal Influences.
Akutsu N., Ooguri M., Onodera T., Kobayashi Y., Katsuyama M., Kunizawa N., Hirao T., Hosoi J., Masuda Y., Yoshida S., Takahashi M., Tsuchiya T. and Tagami H.
Acta Derm Venereol. 89: 21-27, 2009.

Dual Role of COUP-TF-Interacting Protein 2 in Epidermal Homeostasis and Permeability Barrier Formation.
Golonzka O., Liang X., Messaddeq N., Bornert J.-M., Campbell A. L., Metzger D., Chambon P., Ganguli-Indra G., Leid M. and Indra A.K.
J Invest Dermatol. Jun; 129(6): 1459-1470, 2009.

Reservoir based fentanyl transdermal drug delivery systems: effect of patch age on drug release and skin permeation.
Prodduturi S., Smith G.J., Wokovich A.M., Doub W.H., Westenberger B.J. and Buhse L.
Pharm Res. 26:6, 2009.

Novel transdermal delivery of Timolol maleate using sugar esters: Preclinical and clinical studies
El-Laithy H.M.
European Journal of Pharmaceutics and Biopharmaceutic: 72:239–245, 2009.

Transdermal fentanyl in cachectic cancer patients
Heiskanen T., Mätzke S., Haakana S., Gergov M., Vuori E. and Kalso E.
Pain, 144:1:218-222, 2009.

Skin surface electric potential as an indicator of skin condition: a new, non-invasive method to evaluate epidermal condition
Kawau E., Nakanishi J., Kumazawa N., Ozawa K. and Denda M.
Exp Dermatol. 17:688-692, 2008.

Epidermal ceramidase activity regulates epidermal desquamation via stratum corneum acidification
Houben E., Hachem J.P., de Paepe K. and Rogiers V.
Skin Pharmacol Physiol. 21:111-118, 2008.

Increased carbonyl protein levels in the stratum corneum of the face during winter
Kobayashi Y., Iwai I., Akutsu N. and Hirao T.
Int J Cosmet Sci. 30, 35-40, 2008.

In vivo friction study of human skin: influence of moisturizers on different anatomical sites

Ramalho A., Silva C.L., Pais A.A.C.C. and Sousa J.J.S.
Wear 263:1044-1049, 2007.

Cytokines at different stratum corneum levels in normal and sodium lauryl sulphate-irritated skin
de Jongh C.M., Verberk M.M., Spiekstra S.W., Gibbs S. and Kezic S.
Skin Res Technol, Vol 13: 390-398, 2007.

Skin barrier structure and function and their relationship to puritus in end-stage renal disease
Yosipovitch G., Duque M.I., Pate T.S., Ishiuji Y., Guzman-Sanchez A., Dawn A.G., Freedman B.I., Chan Y.H., Crumrine D. and Elias P.M.
Nephrol Dial Transplant, Nov;22(11):3268-72, 2007.

Depth dependence of stratum corneum lipid ordering: a slow-tumbling simulation for electron paramagnetic resonance
Yagi E., Sakamoto K. and Nakagawa K.
J Invest Dermatol, 127:895-899, 2007.

Potential mechanisms by which a single drink of alcohol can increase transdermal absorption of topically applied chemicals
Brand R.M., Jendrzejewski J.L. and Charron A.R.
Toxicology, Jun 25;235(3): 141-149, 2007.

Increased permeability for polyethylene glycols through skin compromised by sodium lauryl sulphate
Jakasa I., Verberk M.M., Bunge A.L., Kruse J. and Kezic S.
Exp Dermatol, Oct;15(10): 801-7, 2006.

Altered penetration of polyethylene glycols into uninvolved skin of atopic dermatitis patients
Jakasa I., Verberk M.M., Esposito M., Bos J.D. and Kezic S.
J Invest Dermatol, 127: 129-134, 2006.

'Keratolytic' properties of benzoyl peroxide and retinoic acid resemble salicylic acid in man
Waller J.M., Dreher F., Benham S., Ford C., Lee C., Tiet T., Weinstein G.D. and Maibach H.I.
Skin Pharmacol Physiol, Vol 19(5): 283-289, 2006.

Percutaneous penetration of sodium lauryl sulphate is increased in uninvolved skin of patients with atopic dermatitis compared with control subjects
Jakasa I., de Jongh C.M., Verberk M.M., Bos J.D. and Kezic S.
Br J Dermatol, Jul;155(1): 104-9 2006.

Variation in barrier impairment and inflammation of human skin as determined by sodium lauryl sulphate penetration rate
De Jongh C.M., Jakasa I., Verberk M.M. and Kezic S.
Br J Dermatol, Apr;154(4): 651-7, 2006.

Transepidermal water loss reflects permeability barrier status: validation in human and rodent in vivo and ex vivo models
Fluhr J.W., Feingold K.R. and Elias P.M.
Exp Dermatol, Jul;15(7): 483-92, 2006.

Determination of polyethylene glycols of different molecular weight in the stratum corneum
Jakasa I., Calkoen F and Kezic S.
J Chromatogr B Analyt Technol Biomed Life Sci. Nov 25; 811(2):177-82, 2004.

Abstracts

Teprenone (Geranylgeranylacetone=GGA), a novel isoprene analog of geranylgeranylphosphate prolongs cell survival and improves skin condition
Lintner K., Mas Chamberlin C., Mondon P and Benech P
Presentation at IFSCC 2007 in Amsterdam, 24-26 Sep 2007.

Variability in the physical properties of the stratum corneum – influences of chronological age and season: 001
Agarwal V., Godfrey M., Long S., Whitby D., Barton S. and Desnos A.
Int J Cosmet Sci, 29(3):219, June 2007.

Mechanism of injury and countermeasures for EVA associated upper extremity medical issues: Extended Vent Tube Study
Jones J., Hoffman R., Harvey G., Bowen C.K., Huday C.E., Tuxhorn J., Gernhardt M. and Scheuring R.
Prenestation at 15th International Academy of Astronautics' "Humans in space" symposium in Beijing, 20-24 May 2007.

Assessing the gentleness of a sonic skin care brush for daily use
Henes E.M., Jackson L., Rodriguez S., Ortblad K. and Akridge R.
J Am Acad Dermatol 2006, 54(3):AB50 Poster 403

Assessment of permeability barrier function measuring transepidermal water loss: comparing 3 closed-loop systems and 4 open-loop systems in vivo and in vitro
Fluhr J. W., Bankova L., Elias P. M. and Feingold K. R.
Skin Res Technol, Vol 9 (2): 164, 2003.

Skin irritation

Influence of tumour necrosis factor- α polymorphism -308 and atopy on irritant contact dermatitis in healthcare workers.
Davis, J.A., Visscher, M.O., Wickett R.R. and Hoath, S. B.
Contact Dermatitis, Dec; 63(6): 320-332, 2010.

Spontaneous itch in the absence of hyperalgesia in a mouse hindpaw dry skin model.
Akiyama T., Carstens M.I. and Carstens E.
Neuroscience Letters, Oct 22; 484(1), 62-75, 2010.

A double-blind, randomized study to assess the effectiveness of different moisturizers in preventing dermatitis induced by hand washing to simulate healthcare use.
Williams C., Wilkinson S., McShane P., Lewis J., Pennington D., Pierce S. and Fernandez C.
Br J Dermatol, May; 162(5): 1088-1092, 2010.

Effects of glycerol on human skin damaged by acute sodium lauryl sulphate treatment.
 Atrux-Tallau N., Romagny C., Padois K., Denis A., Haftek M., Falson F., Pirot F. and Maibach H.I.
 Arch Dermatol Res, 302: 435-441, 2010.

Role of TNF-polymorphism -308 in irritant contact dermatitis and neurosensory response.
 Davis J.A.
 PhD Thesis. Department of Pharmaceutical Sciences of the College of Pharmacy, University of Cincinnati, 2009.

The alkaline pH-adapted skin barrier is disrupted severely by SLS-induced irritation
 Kim E., Kim S., Nam G.V., Lee H., Moon S and Chang I
 Int J Cosmet Sci, 31:4:263-269, 2009.

Skin Barrier Disruption by Sodium Lauryl Sulfate-Exposure Alters the Expressions of Involucrin, Transglutaminase 1, Profilaggrin, and Kallikreins during the Repair Phase in Human Skin In Vivo.
 Törmä H., Lindberg M. and Berne B.
 J Invest Dermatol, epub 15 Nov 2007.

Differential cytokine expression in skin after single and repeated irritation by sodium lauryl sulphate
 de Jongh C.M., Lutter R., Verberk M.M. and Kezic S.
 Exp Dermatol, 16, 1032-1040, 2007.

A randomized, controlled, double-blind study of the effect of wearing coated pH 5.5 latex gloves compared with standard powder-free latex gloves on skin pH, transepidermal water loss and skin irritation.
 Mirza R., Maani N., Liu C., Kim J. and Rehmus W.
 Contact Dermatitis, Jul;55(1): 20-5, 2006.

Stratum corneum cytokines and skin irritation response to sodium lauryl sulfate
 de Jongh C.M., Verberk M.M., Withagen C.E.T., Jacobs J.J.L., Rustemeyer T. and Kezic S.
 Contact Dermatitis, Jun;54(6):325-33, 2006.

Skin reaction and recovery: a repeated sodium lauryl sulphate patch test vs. a 24-h patch test and tape stripping
 Koopman D.G., Kezic S., Verberk M.M.
 Br J Dermatol, Mar;150(3):493-9, 2004.

Effect of skin care, treatment, nutrition

Influence of tumour necrosis factor- α polymorphism –308 and atopy on irritant contact dermatitis in healthcare workers.
 Davis, J.A., Visscher, M.O., Wickett R.R. and Hoath, S. B.
 Contact Dermatitis, Dec; 63(6): 320–332, 2010.

Tolerance of Baby Cleansers in Infants: A Randomized Controlled Trial.
 Dizon M.V., Golzote C., EstanislaoR., Mathew N. and Sarkar R.
 Indian Pediatrics, Nov 7; 47(11): 959-63, 2010.

Effect of Oral Intake of Ceramide-Containing Acetic Acid Bacteria on Skin Barrier Function.
 Oda T., Tachimoto H., Kishi M. and Kaga T. and Ichihashi M.
 Anti-Aging Medicine, 7(5), 50-54, 2010. Published online: Mar. 25, 2010

5-Aminolevulinic Acid-based Photodynamic Intense Pulsed Light Therapy Shows Better Effects in the Treatment of Skin Photoaging in Asian Skin.
 Yang G., Xiang L.H. and Gold M.H.
 J Clin Aesthet Dermatol., March; 3(3): 40-43, 2010.

A randomized, controlled comparative study of the wrinkle reduction benefits of a cosmetic niacinamide/peptide/retinyl propionate product regimen vs. a prescription 0,02% tretinoin product regimen.
 Fu J., Hillebrand G., Raleigh P., Li J., Marmor M., Bertucci V., Grimes P., Mandy S., Perez M., Weinkle S. and Kaczvinsky J.
 Br J Dermatol, March; 162(3): 647–654, 2010.

The Safety and Efficacy of Pimecrolimus, 1%, Cream for the Treatment of Netherton Syndrome.
 Yan A.C., Honig P.J., Ming M.E., Weber J. and Shah K.N.
 Arch Dermatol., 146(1): 57-62, 2010.

A novel organogermanium protected atopic dermatitis induced by oxazolone.
 Lim D.H., Li M., Seo J.-A., Lim K.-M. and Ham S.W.
 Bioorg Med Chem Lett, Jul 15; 20(14): 4032-4034, 2010.

A double-blind, randomized study to assess the effectiveness of different moisturizers in preventing dermatitis induced by hand washing to simulate healthcare use.
 Williams C., Wilkinson S., McShane P., Lewis J., Pennington D., Pierce S. and Fernandez C.
 Br J Dermatol, May; 162(5): 1088–1092, 2010.

Comparison of skin barrier function and sensory nerve electric current perception threshold between IgE-high extrinsic and IgE-normal intrinsic types of atopic dermatitis.
 Mori T., Ishida K., Mukumoto S., Yamada Y., Imokawa G., Kabashima K., Kobayashi M., Bito T., Nakamura M., Ogasawara K. and Tokura Y.
 Br J Dermatol, Jan; 162(1): 83–90, 2010.

Skin Care in the NICU Patient: Effects of Wipes versus Cloth and Water on Stratum Corneum Integrity.
 Visscher M., Odio M., Taylor T., White T., Sargent S., Sluder L., Smith L., FlowerT., Mason B., Rider M., Huebner A. and Bondurant P.
 Neonatology. 96(4): 226-34, 2009.

Glycylglycine decreases the size of conspicuous facial pores: Single-blinded half areas of face-applied study.
 Iida T., Kaneko M., Kawai E., Muta-Takada K. and Inomata S.
 J Dermatol. Feb; 36(2): 120-123, 2009.

Role of TNF-polymorphism -308 in irritant contact dermatitis and neurosensory response.
 Davis J.A.

PhD Thesis. Department of Pharmaceutical Sciences of the College of Pharmacy, University of Cincinnati, 2009.

Cosmeceutical properties produced by levan produced by Zymomonan mobilis
Kim K.H., Chung C.B., Kim Y.H., Kim K.S., Han C.S. and Kim C.H.
J Cosmet Sci, 56:395-406, 2005.

Evaluation of the antioxidant capacity and preventive effects of a topical emulsion and its vehicle control on the skin response to UV exposure
Zhai H., Behnam S., Villarama C.D., Arens-Corell M., Choi M.J. and Maibach H.I.
Skin Pharmacol Physiol, 18: 288-293, 2005.

Efficacy of dietary hempseed oil in patients with atopic dermatitis
Callaway J., Schwab U., Harvima I., Halonen P., Mykkänen O., Hyvönen P. and Järvinen T.
Dermatology Treat, 16: 87-94, 2005.

Moisturizing effects of topical nicotinamide on atopic dry skin
Soma Y., Kashima M., Imaizumi A., Takahama H., Kawakami T. and Mizoguchi M.
Int J Dermatol, Mar;44(3): 197-202, 2005.

Abstracts

Topical coconut oil application reduces trans-epidermal water loss (TEWL) in preterm very low birth weight neonates: A randomised clinical trial

Nangia S., Paul V.K. and Deorari A.K.
48th annual meeting of ESPR in Prague, Check Republic, 6 –8 Oct 2007, p. 105.
Pediatrics, Vol 121 Suppl 2, Jan 2008.

A pilot study of the efficacy and safety of pimecrolimus 1% cream in the treatment of Netherton syndrome: Interim results
Yan A., Shah K., Kubrak C. and Parneix-Spake A.
J Am Acad Dermatol, February: AB153, 2007.

Susceptibility to SLS irritation: Effect of a five-day pre-use of two skin care oils studied comparing various techniques and instruments
Rantanen, T., Snellman, E. and Kostainen M.
First World Congress on Work-Related and Environmental Allergy, 9-12 July 2003: p. 95.

Scalp, lip and nail measurements

Seasonal factors influencing biophysical properties of facial and labial skin.
Gubanova E.I., Glad'ko V.V., Cazei L., Rodina M.Yu. and Lapatina N.G.
Klin Dermatol Venerol, 2:19-24, 2010. Russian.

Tape stripping on a human nail: quantification of removal
Tudela E., Lamberbourg A., Diaz A.C., Zhai H. and Maibach H.I.
Skin Res Technol, 14: 472-477, 2008.

The use of an extended ventilation tube as a countermeasure for EVA-associated upper extremity medical issues
Jones J.A., Hoffman R.B., Buckland D.A., Harvey C.M., Bowen C.K., Hudy C.E. Strauss S., Novak J. and Gernhardt M.L.
Acta Astronaut, 63:763-768, 2008.

Influence of age and hormone replacement therapy on the functional properties of the lips
Caisey L., Gubanova E., Camus C., Lapatina N., Smetnik V. and Lévéque J.-L.
Skin Res Technol, 14:220-225, 2008.

Water loss through lip, nail, eyelid skin, scalp skin and axillar skin measured with a closed-chamber evaporation principle
Nuutilinen J., Harvima I., Lahtinen M.-R. and Lahtinen T.
Br J Dermatol, 148: 839-841, 2003.

Abstracts

Shampooing with pyrithione zinc reduces transepidermal water loss in scalp of dandruff-involved patients
Billhimer W., Erb J., Bacon R., Ertel K and Copas M.
J Am Acad Dermatol 2006, 54(3):Poster

Comparison of two instruments to assess scalp barrier function
Bacon R., Billhimer, W., Ertel, K. and Watson, D.
Skin Res Technol, Vol 10: 13, 2004.

Wound healing

Epidermal wound healing in severe sepsis and septic shock in humans.
Koskela M., Gäddnäs F., Ala-Kokko T.I., Laurila J.J., Saarnio J., Oikarinen A. and Koivukangas V.
Crit Care. Jun; 13(3): R100, 2009.

Instrumental evaluation of the protective effects of a barrier film on surrounding skin in chronic wounds
Dini V., Salibra F., Brilli C. and Romanelli M.
Wounds, 20:254-257, Sept 2008.

Use of nanotechnology-designed footsock in the management of preulcerative conditions in the diabetic foot: results of a single, blind randomized study
Banchellini E., Macchiarini S., Dini V., Rizzo L., Tedeschi A., Scatena A., Goretti C., Campi F., Romanelli M.. and Alberto P
Int J Low Extrem Wounds, 7:2:82-87, 2008.

Abstracts

Moisture vapour transmission rate – How is this affected by wound fluid viscosity?
Adams S., Pritchard D. and Rogers J.

Evaluation of maceration by trans epidermal water loss (TEWL) in patients with venous leg ulcers
Dini V., Salibra F., Brilli C., Bertone M. and Romanelli M.
Sumposium on Advanced Wound Care & Medical Research Forum on Wound Repair, p. 128, 2006.

Skin surface water loss SSWL

Abstracts

Measurement of skin hydration in the diapered area of infants with a closed chamber TEWL meter
Alanen, E., Wiesemann, F., Nuutinen, J. and Lahtinen, T.
Skin Res Technol, Vol 9 (2): 199, 2003.

Physiological and Sympathetic skin response, Hyperhidrosis

Objective evaluation of patients with palmar hyperhidrosis submitted to two levels of sympathectomy: T3 and T4.
Ishy A., de Compos J.R.M., Wolosker N., Kauffman P., Tedde M.L., Chiavoni A.R. and Janete F.B.
Interact Cardiovasc Thorac Surg, published online Jan 13 2011.

Effect of Itch, Scratching and Mental Stress on Autonomic Nervous System Function in Atopic Dermatitis.
Tran B.W., Papouli A.D.P., Russoniello C.V., Wang H., Patel T.S.; Chan Y.-H. and Yosipovitch G.
Acta Derm Venereol, Jun; 90(4), 354-361, 2010.

Endoscopic lumbar sympathectomy for plantar hyperhidrosis.
Rieger R., Pedevilla S. and Pöchlauer S.
Br J Surg. Dec; 96(12): 1422-1428, 2009.

Effects of low humidity and high air velocity in a heated room on physiological responses and thermal comfort after bathing:
An experimental study
Hashiguchi N. and Tochihara Y.
Int J Nurs Stud 46:172-180, 2009.

Primary Palmoplantar Hyperhidrosis and Thoracoscopic Sympathectomy: A New Objective Assessment Method
Tetteh H.A., Groth S.S., Kast T., Whitson B.A., Radosevich D.M., Klopp A.C., D'Cunha J., Maddaus M.A. and Andrade R.S.
Ann Thorac Surg, 87:1:267-275, January 2009.

Increased carbonyl protein levels in the stratum corneum of the face during winter
Kobayashi Y., Iwai I., Akutsu N. and Hirao T.
Int J Cosmet Sci, 30:35-40, 2008.

Endoscopic lumbar sympathectomy for women: Effect on compensatory sweat
Loureiro M.P., de Campos J.R.M., Kauffman P., Jatene F.B., Weigmann S. and Fontana A.
Clinics, 63(2):189-96, 2008.

Upper limb musculoskeletal disorders - With special reference to sympathetic nerve functions and tactile sensation
Ming Z.
Kuopio University Publications D. Medical Sciences 405. 2007. 91 p. MD Thesis.

Retroperitoneoscopic lumbar sympathectomy for the treatment of plantar hyperhidrosis: technique and preliminary findings
Rieger R. and Pedevilla S.
Surg Endosc, Jan;21(1):129-35, 2007.

Physiological and subjective responses to low relative humidity in young and elderly men
Sunwoo Y., Chou C., Takeshita J., Murakami M. and Tochihara Y.
J Physiol Anthropol, 25:229-238, 2006.

Physiological and subjective responses to low relative humidity
Sunwoo Y., Chou C., Takeshita J., Murakami M. and Tochihara Y.
J Physiol Anthropol, 25:7-14, 2006.

Abstracts

Objective evaluation of thoracoscopic sympathectomy for palmar hyperhidrosis
Tetteh H.A., Kast T., Maddaus M.M., Whitson B.A., Groth S.S., Radosevich D.M. and Andrade R.S.
44th Annual Meeting, The Society of Thoracic Surgeons, Jan 28-30, 2008, p. 176

Measurement of the amount of average hydrosis for giving data for scientific comparison
Pedevilla S. and Rieger R.
Presented in 7th International Symposium on Sympathetic Surgery, 20-22 March 2007
Clinical Autonomic Research, Vol 17, Number 2 / April 2007.

Veterinary dermatology and skin behaviour, treatment and nutrition, Zoology

Spontaneous itch in the absence of hyperalgesia in a mouse hindpaw dry skin model.
Akiyama T., Carstens M.I. and Carstens E.
Neuroscience Letters, Oct 22; 484(1), 62-75, 2010.

Metabolic adaptations to heat stress in growing cattle.
O'Brien M.D., Rhoads R.P., Sanders S.R., Duff G.C. and Baumgard L.H.
Domestic Animal Endocrinology, Feb; 38(2), 86-94, 2010.

Induced keratinocyte hyper-proliferation in $\alpha 2\beta 1$ integrin transgenic mice results in systemic immune cell activation.
Teige I., Backlund A., Svensson L., Kvist P.H., Petersen T.K. and Kemp K.

International Immunopharmacology, Jan; 10(1), 2010.

Transepidermal Water Loss, Stratum Corneum Hydration and Transmission Electron Microscope Observation on Acetone Damaged Canine Skin Barrier Model.
Oh W., Park S. and Oh T.
J Vet Clin, 27(2): 142-146, (2010).

Mapping of the dog skin based on biophysical measurements.
Oh W.-S. and Oh T.-H.
Vet Derm, 21: 367-372, 2010.

Evaluation of a hand-held evaporimeter (VapoMeter[®]) for the measurement of transepidermal water loss in healthy dogs.
Lau-Gillard P. J., Hill P. B., Chesney C. J., Budleigh C. and Immonen A.
Vet Derm, 21: 136-145, 2010.

Evaluation of physiological and blood serum differences in heat-tolerant (Romosinuano) and heat-susceptible (Angus) Bos taurus cattle during controlled heat challenge.
Scharf B., Carroll J.A., Riley D.G., Chase C.C. Jr., Coleman S.W., Keisler D.H., Weaver R.L. and Spiers D.E.
J Anim Sci. Jul; 88(7): 2321-36, 2010.

Auswirkung von lokal applizierten Fettsäuren auf den epidermalen Wasserverlust, Pruritus und klinischen Hautzustand atopischer und gesunder Hunde.
Tretter S.
PhD Thesis, Ludwig-Maximilians-Universität München, 2010.

Measurement of transepidermal water loss from clipped and unclipped anatomical sites on the dog.
Oh W.-S. and Oh T.-H.
Aust Vet J. Oct; 87(10): 409-412, 2009.

Effects of a supplemental yeast culture on heat-stressed lactating Holstein cows.
Shwartz G., Rhoads M.L., VanBaale M.J., Rhoads R.P. and Baumgard L.H.
J Dairy Sci. Mar; 92(3): 935-942, 2009.

A Re-Evaluation of the Impact of Temperature Humidity Index (THI) and Black Globe Humidity Index (BGHI) on Milk Production in High Producing Dairy Cows.
Zimbelman R. B., Rhoads R.P., Rhoads M.L., Duff G.C., Baumgard L. H. and Collier R. J. Southwest Nutrition & Management Conference, Arizona, Feb 26.-27. 2009.

Water exchange and permeability properties of skin in three species of amphibious sea snakes (*Laticauda* spp.).
Lillywhite H.B., Menon J.G., Menon G.K., Sheehy 3rd C.M. and Tu M.C.
J Exp Biol, 212:1921-1929, 2009.

Profile of the bovine acute-phase response following an intravenous bolus-dose lipopolysaccharide challenge
Carroll J.A., Reuter R.R., Chase Jr C.C., Coleman S.W., Riley D.G., Spiers D.E., Arthington J.D. and Galyean M.L.
Innate Immunity, 15:2:81-89, 2009.

Chronic ethanol ingestion alters xenobiotic absorption through the skin: Potential role of oxidative stress
Brand R.M. and Jendrzejewski J.L.
Food Chem Toxicol, 46:1940-1948, 2008.

Olopatadine ameliorates rat experimental cutaneous inflammation by improving skin barrier function
Tamura T., Matsubara M., Amano T. and Chida M.
Pharmacology; 81:118-126, 2008.

Potential mechanisms by which a single drink of alcohol can increase transdermal absorption of topically applied chemicals
Brand R.M., Jendrzejewski J.L. and Charron A.R.
Toxicology, Jun 25;235(3): 141-149, 2007.

Olopatadine hydrochloride accelerates the recovery of skin barrier function in mice
Amano T., Takeda T., Yano H. and Tamura T.
Br J Dermatol, May;156(5):906-912, 2007.

Cutaneous water loss and lipids of the stratum corneum in dusky anticirds, a lowland tropical bird
Munos-Garcia A. and Williams J.B.
The Condor, 109:59-66, 2007.

Transepidermal water loss reflects permeability barrier status: validation in human and rodent in vivo and ex vivo models
Fluhr J.W., Feingold K.R. and Elias P.M.
Exp Dermatol, Jul;15(7): 483-92, 2006.

Abstracts

Skin permeability and water relations of sea snakes
Lillywhite H.B., Babonis L.S., Shhehy C.M. and Tu M.-C.
FASEB J, 22:1239.18, 2008.

Sweating rate of dairy cows under shade and sunny environments
Gebremedhin K.G., Hillman P.E., Lee C.N. and Collier R.J.
Presented at American Society of Agricultural and Biological Engineers Annual Meeting, 2007.

Regional differences in sweat rate response of steers to short-term heat stress
Spiers D.E., Wax L.E., Scharf B. and Aiken G.E.
J Anim Sci, Vol 84, Suppl 1/J Dairy Sci, Vol 89, Suppl 1: 303-304, 2006.

Effects of ambient temperature and solar radiation on skin evaporative water loss in dairy cattle
Pollar B.C., Dwyer, M.E., Fitzgerald A.C., Gentry P.C., Henderson D.A. and Collier R.J.
J Anim Sci, Vol 82, Suppl 1/J Dairy Sci, Vol 87, Suppl 1/Poult Sci Vol 83, Suppl 1: 198, 2006.

Impact of alcohol consumption on the skin as determined by non-invasive bioengineering tools

Effects of ambient temperature and solar radiation on skin evaporative water loss in dairy cattle
Pollard, B. C., Dwyer, M. E., Fitzgerald, A. C., Gentry, P. C., Henderson, D. A. and Collier, R. J.
J Anim Sci, Vol 82, Suppl. 1/ *J Dairy Sci*, Vol 87, Suppl. 1/ *Poult Sci*, Vol 83, Suppl. 1. 198, 2004.

Heat dissipation in winter-acclimated lactating cows and non-lactating heifers subjected to increased ambient temperature and solar radiation in Arizona
Pollard, B. C., Annen, E. A., Baumgard, L. H., Cheatham, R. C., Estheimer, M. D., Dwyer, M. E., Fitzgerald, A. C., Halfliger, H. C., Moore, C. E., Kay, J. K., Mendivil, O. B., Gentry, P. C., Henderson, D. A., Steining, C. M. and Collier, R. J.
J Anim Sci, Vol 82, Suppl. 1/ *J Dairy Sci*, Vol 87, Suppl. 1/ *Poult Sci*, Vol 83, Suppl. 1. 461, 2004.

In-vitro studies

Magnetophoresis for enhancing transdermal drug delivery: Mechanistic studies and patch design.
Murthy, S.N., Sammeta S.M. and Bowers C.
Journal of Controlled Release, Dec 1; 148(2): 197-203, 2010.

Evaluation needle length and density of microneedle arrays in the pretreatment of skin for transdermal drug delivery.
Yan G., Warner K.S., Zhang J., Sharma S. and Gale B.K.
Int J Pharm, May 31; 391(1-2): 7-12, 2010.

Microchannels created by sugar and metal microneedles: Characterization by microscopy, macromolecular flux and other techniques.
Li G., Badkar A., Kalluri H. and Banga A. K.
J Pharm Sci. Apr; 99(4): 1931-41, 2010.

Transdermal iontophoretic delivery of terbinafine hydrochloride: Quantitation of drug levels in stratum corneum and underlying skin.
Sachdeva V., Siddaju S., Yu Y.-Y., Kim H.D., Friden P.M. and Banga A.K.
Int J Pharm, Mar; 388(1-2), 24-31, 2010.

Transdermal and intradermal iontophoretic delivery of dexamethasone sodium phosphate: quantification of the drug localized in skin.
Paturi J., Kim H.D., Chakraborty B., Friden P.M. and Banga A.K.
J Drug Targ, Feb; 18(2): 134-140, 2010.

Transdermal delivery of fentanyl from matrix and reservoir systems: Effect of heat and compromised skin.
Prodduturi S., Sadrieh N., Wokovich A.M., Doub W.H., Westenberger B. J. and Buhse L.
J Pharm Sci, 99: 2357-2366, 2010.

Microneedle Arrays Allow Lower Microbial Penetration Than Hypodermic Needles In Vitro.
Donnelly R.F., Singh T.R.R., Tunney M.M., Morrow D.I.J., McCarron P.A., O'Mahony C. and Woolfson A.D.
Pharm Res. Nov; 26(11): 2513-2522, 2009.

Poly(amidoamine) dendrimers as skin penetration enhancers: Influence of charge, generation, and concentration.
Venuganti V.V.K. and Perumal O.P.
J Pharm Sci, Jul; 98(7): 2345-2356, 2009.

OcuDrain-E™ -A noninvasive technique for reduction of intraocular pressure.
Murthy S.M., Sammeta S.M. and Vaka S.R.K.
Int J Pharm, Mar; 369(1-2): 92-95, 2009.

Skin penetration enhancement by a microneedle device (Dermaroller®) in vitro: Dependency on needle size and applied formulation.
Badran M.M., Kuntsche J. and Fahr A.
Eur J Pharm Sci. Mar; 36(4-5): 511-523, 2009.

Reservoir based fentanyl transdermal drug delivery systems: effect of patch age on drug release and skin permeation.
Prodduturi S., Smith G.J., Wokovich A.M., Doub W.H., Westenberger B.J. and Buhse L.
Pharm Res, 26:6, 2009.

Molecular charge mediated transport of a 13 kD protein across microporated skin
Katikaneni S., Badkar A., Nema S and Banga A.K.
Int J Pharm, 378:93-100, 2009.

Films based on chitosan polyelectrolyte complexes for skin drug delivery:Development and characterization
Silva C.L., Pereira J.C., Ramalho A., Pais A.C.C.C., Sousa J.J.S.
J Memb Sci, 320:268-279, 2008.

Effect of poly(amidoamine) (PAMAM) dendrimer on skin permeation of 5-fluorouracil
Venuganti V.V.K. and Perumal O.P.
Int J Pharm, 361:230-238, 2008.

Interaction of lipid nanoparticles with human epidermis and an organotypic cell culture model
Kuntsche J., Bunjes H., Fahr A., Pappinen S., Rönkkö S., Suhonen M. and Urtti A.
Int J Pharm, 354:180-195, 2008.

Qualitative and quantitative comparison of heat separated epidermis and dermatomed skin in percutaneous absorption studies
Atrux-Tallau N., Pirot F., Falson F., Roberts M.S. and Maibach H.I.
Arch Dermatol Res, 299:507-511, 2007.

In vitro human skin model to evaluate water permeability and determine wound dressings' occlusivity
Zhai X., Yokota M. and Maibach H.I.
Cutan Ocul Toxicol ;26(2):107-11, 2007.

Vitamin C enhances differentiation of a continuous keratinocyte cell line (REK) into epidermis with normal stratum corneum ultrastructure and functional permeability barrier
Pasanen-Seppänen S., Suhonen T.M., Kirjavainen M., Suihko E., Urtti A., Miettinen M., Hyttinen M., Tammi M. and Tammi R.
Histocem Cell Biol, 116: 287-297, 2001.

Formation of permeability barrier in epidermal organotypic culture for studies on drug transport
Pasonen-Seppänen S., Suhonen T.M., Kirjavainen M., Miettinen M., Urtti A., Tammi M. and Tammi R.
J Invest Dermatol, Vol 117, No 5, 1322-1324, November 2001.

Validation articles

Tape-stripping method in man: comparison of evaporimetric methods
Zhai H., Dika E., Goldovsky M. and Maibach H.I.
Skin Res Technol, Vol 13: 207-210, 2007.

Measurement of transepidermal water loss by closed-chamber systems
Nuutinen J.
in Serup J., Jemec G.B.E. and Grove G., (eds.), Handbook of non-invasive methods and the skin, 2nd edition, CRC Press, Boca Raton, FL, pp. 393-396, 2006.

Comparative evaporimetry in man
Shah J.H., Zhai H. and Maibach H.I.
Skin Res Technol, Vol 11: 205-208, 2005.

Validation of the VapoMeter, a closed unventilated chamber system to assess transepidermal water loss vs. the open chamber Tewameter
De Paepe K., Houben E., Adam R., Wiesmann F. and Rogiers V.
Skin Res Technol, Vol 11: 61-69, 2005.

Water loss through lip, nail, eyelid skin, scalp skin and axillar skin measured with a closed-chamber evaporation principle
Nuutinen J., Harvima I., Lahtinen M.-R. and Lahtinen T.
Br J Dermatol, 148: 839-841, 2003.

A closed unventilated chamber for the measurement of transepidermal water loss
Nuutinen J., Alanen E., Autio P., Lahtinen M-R., Harvima I. and Lahtinen T.
Skin Res Technol, Vol 9: 85-89, 2003.

Abstracts

Applicability of the closed unventilated VapoMeter to assess transepidermal water loss (TEWL) on nasolabial skin areas
De Paepe, K., Houben, E., Adam, R., Wiesemann, F. and Rogiers, V.
Skin Res Technol, Vol 10: 3, 2004.

Comparison of two methods for TEWL measurement
Shah J.H., Zhai H., Maibach H.I. and Nuutinen J.
Skin Res Technol, Vol 10, 2004.

Measurement of water evaporation rates with a closed chamber TEWL meter
Nuutinen, J., Alanen, E. and Lahtinen, T.
Skin Res Technol, Vol 9 (2): 183, 2003.

MoistureMeter D

Lymphedema

Randomised phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphoedema after radiotherapy for cancer.
 Gothard L., Haviland J., Bryson P., Laden G., GloverM., HarrisonS., Woods M., Cook G., Peckitt C., Pearson A., Somaiah N., Stanton A., Mortimer P. and Yarnold J.
Radiother Oncol. Oct; 97(1): 101-107, 2010.

Local Tissue Water in At-Risk and Contralateral Forearms of Women with and without Breast Cancer Treatment-Related Lymphedema.
 Mayrovitz H.M., Weingrad D.N. and Davey S.
Lymphat Res Biol. Sep; 7(3): 153-158, 2009.

Assessing lymphedema by tissue indentation force and local tissue water.
 Mayrovitz H.N.
Lymphology, Jun; 42(2): 88-98, 2009.

Suitability of single tissue dielectric constant measurements to assess local tissue water in normal and lymphedematous skin
 Mayrovitz H.N., Davey S. and Shapiro E.
Clin Physiol Imaging, 29(2):123-127, 2009.

Local tissue water assessed by tissue dielectric constant: anatomical site and depth dependence in women prior to breast cancer treatment-related surgery.
 Mayrovitz H.N., Davey S. and Shapiro E.
Clin Physiol Imaging, 28:337-342, 2008.

Localized tissue water changes accompanying one manual lymphatic drainage (MLD) therapy session assessed by changes in tissue dielectric constant inpatients with lower extremity lymphedema.
 Mayrovitz H.N., Davey S. and Shapiro E.
Lymphology, 41:87-92, 2008.

Assessing local tissue edema in postmastectomy lymphedema.
 Mayrovitz H.N.
Lymphology, 40:87-94, 2007.

A new water-specific technique for local measurement of edema and lymphedema induced by cancer therapy
 Lahtinen T., Nuutinen J. and Mayrovitz H.N.
ESTRO Newsletter 61: 23-24, Winter 2006.

Abstracts

A novel MoistureMeter D for local assessment of lymphedema
 Niinimäki M., Nuutinen J. and Lahtinen T.
The European Journal of Lymphology and related problems, 20:57:13, 2009.

Local tissue water changes accompanying a single manual lymphatic drainage (MLD) treatment in patients with lower extremity lymphedema
 Mayrovitz H.N., Davey S. and Shapiro E.
 Poster presented at 8th National Lymphedema Network (NLN) International Conference, August 2008.

Local tissue water assessed by tissue dielectric constant: anatomical site and depth dependence in women prior to breast cancer related surgery
 Mayrovitz H.N., Davey S. and Shapiro E.
 Poster presented at 8th National Lymphedema Network (NLN) International Conference, August 2008.

Assessment of local tissue edema in arms of women with postmastectomy lymphedema
 Mayrovitz H.N. and Brown-Cross D.
2006 Symposium on Advanced Wound Care & Medical Research Forum on Wound Repair, p.189

Cutaneous edema, Tissue fluid status

Male-female differences in forearm skin tissue dielectric constant.
 Mayrovitz H. N., Carson S. and Luis M.
Clin Physiol Funct Imaging. Sep; 30(5): 328-32, 2010.

5-Aminolevulinic Acid-based Photodynamic Intense Pulsed Light Therapy Shows Better Effects in the Treatment of Skin Photoaging in Asian Skin.
 Yang G., Xiang L.H. and Gold M.H.
J Clin Aesthet Dermatol., Mar; 3(3): 40-43, 2010.

Noninvasive transcutaneous sampling of glucose by electroporation
 Murthy S.S., Kiran V.S.R., Mathur S.K. and Murthy S.N.
J Diabetes Sci Technol, 2(3):250-254, 2008.

Chronic ethanol ingestion alters xenobiotic absorption through the skin: Potential role of oxidative stress
 Brand R.M. and Jendrzejewski J.L.
Food Chem Toxicol, 46:1940-1948, 2008.

Moisturizing lotions can increase transdermal absorption of the herbicide 2,4-dichlorophenoxyacetic acid across hairless mouse skin
 Brand R.M., Charron A.R., Sandler V.L. and Jendrzejewski J.L.
Cutan Ocul Toxicol, 26:15-23, 2007.

Potential mechanisms by which a single drink of alcohol can increase transdermal absorption of topically applied chemicals
 Brand R.M., Jendrzejewski J.L. and Charron A.R.
Toxicology, Jun 25;235(3): 141-149, 2007.

Skin tissue water and laser Doppler blood flow during a menstrual cycle
Mayrovitz H.N., Brow-Cross D. and Washington Z.
Clin Physiol Funct Imaging, 27: 54-59, 2007.

Measurement of oedema in irritant-exposed skin by a dielectric technique
Miettinen M., Mönkkönen J., Lahtinen M.R., Nuutinen J. and Lahtinen T.
Skin Res Technol, Vol 12: 235-240, 2006.

Validation of a new dielectric device to assess changes of tissue water in skin and subcutaneous fat
Nuutinen J., Ikäheimo R. and Lahtinen T.
Physiol Meas, 25: 447-454, 2004.

Dielectric constant of skin and subcutaneous fat to assess fluid changes after cardiac surgery
Petäjä L., Nuutinen J., Uusaro A., Lahtinen T. and Ruokonen E.
Physiol Meas, 24: 383-390, 2003.

Quantitative assessment of protein content in irradiated human skin
Lahtinen T., Nuutinen J., Alanen E., Turunen M., Nuortio L., Usenius T. and Hopewell J.W.
Int J Radiat Oncol Biol Phys, 43(3): 635-638, 1999.

Abstracts

Impact of alcohol consumption on the skin as determined by non-invasive bioengineering tools
Brand R.M., Jendrzejewski J.L. and Charron A.R.
Society of Toxicology Meeting, March 2006.

Water content at skin surface and in whole skin during irritant skin reaction
Miettinen M., Nuutinen J., Lahtinen M.-R. and Lahtinen T.
13th Congress of European Academy of Dermatology and Venereology Meeting, 17-21 November, 2004.

Measurement of skin edema by a dielectric technique (MoistureMeter-D)
Miettinen M., Lahtinen M.-R., Nuutinen J., Lahtinen T. and Niinimäki M.
Contact Dermatitis, 50:18, 2004.

EdemaMeter - a new device to detect tissue edema in skin and subcutaneous tissues
Lahtinen T., Ikäheimo R., Lampainen E. and Nuutinen, J.
Skin Res Technol, Vol, 9 (2): 182, 2003.

Radiotherapy

A dielectric method for measuring early and late reactions in irradiated human skin
Nuutinen J., Lahtinen T., Turunen M., Alanen E., Tenhunen M., Usenius T. and Kolle R.
Radiother Oncol, 47: 249-254, 1998.

Skin dielectric constant at high radiofrequency with special emphasis on radiation-induced late skin reaction
Nuutinen, J.
Kuopio University Publications C. Natural and Environmental Sciences 55. 1997. 64 p. Ph.D. Thesis.

Burns, Thermal injury

Dielectric measurement in experimental burns: A new tool for burn depth determination?
Soo C. and Garner W.L.
Plast Reconstr Surg, Mar;117(3): 899-901, 2006.

Dielectric measurement in experimental burns: a new tool for burn depth determination?
Papp A., Lahtinen T., Harma M., Nuutinen J., Uusaro A. and Alhava E.
Plast Reconstr Surg, Mar;117(3): 889-98; discussion 899-901, 2006.

Experimental thermal injury: New methods in assessing tissue damage
Papp A.
Kuopio University Publications D. Medical Sciences 356. 2005. 78 p. MD Thesis

Wound healing

Abstracts

Assessing a Biophysical Device to Predict Pressure Ulcers in Spinal Cord Injury
Bates-Jensen B., Guihan M. McCreath H., Chun S., Parachuri R and Simon J.
Poster presented at 2nd International Evidence Practice in Wound Care, USA, June 2009.

Initial assessment of tissue water content surrounding pressure ulcers in spinal cord injury patients
Harrow J.J. and Mayrovitz H.N.
2006 Symposium on Advanced Wound Care & Medical Research Forum on Wound Repair, p.73

Weight loss

Changes in abdominal subcutaneous fat water content with rapid weight loss and long-term weight maintenance in abdominally obese men and women
Laaksonen D.E., Nuutinen J., Lahtinen T., Rissanen A., and Niskanen L.K.
Int J Obesity, 27: 677-683, 2003.

Theoretical background

Male-female differences in forearm skin tissue dielectric constant.
 Mayrovitz H. N., Carson S. and Luis M.
Clin Physiol Funct Imaging. Sep; 30(5): 328-32, 2010.

Dielectric properties of the skin
 Lahtinen, T., Nuutinen, J., Alanen, E.
 In: Radio Frequency Radiation Dosimetry. Editors: B. J. Klauenberg and D. Miklavcic. Kluwer Academic Publishers, the Netherlands, 2000

Penetration of electromagnetic fields of open-ended coaxial probe between 1 MHz and 1GHz in dielectric skin measurements
 Alanen, E., Lahtinen T. and Nuutinen J.
Phys Med Biol, 44(7): N169-176, 1999.

Variational formulation of open-ended coaxial line in contact with layered biological medium
 Alanen, E. Lahtinen, T. and Nuutinen, J.
IEEE Trans Biomed Eng, 45 (10) 1241-1248, 1998.

Measurement of dielectric properties of subcutaneous fat with open-ended coaxial sensors
 Alanen, E., Lahtinen, T. and Nuutinen, J.
Phys Med Biol, 43: 475-485, 1998.

Dielectric properties of the skin
 Lahtinen, T., Nuutinen, J. and Alanen, E.
Phys Med Biol, 42: 1471-1472, 1997.

Abstracts

Measurement depths of a skin-water analyzer (MoistureMeter D)
 Kiiskinen M., Nuutinen J. and Alanen E.
Skin Res Technol, Vol 11: 292, 2005.

MoistureMeter SC

Strontium hexaferrite nanomagnets suspended in a cosmetic preparation: a convenient tool to evaluate the biological effects of surface magnetism on human skin.
 Zioni T., Perkas N., Wolfus Y., Soroka Y., Popov I., Oron M., Perelshtein I., Bruckental Y., Brégégère F. M., Ma'or Z., Gedanken A., Yeshurun Y., Neuman R. and Milner Y.
Skin Res Technol. Aug; 16(3): 316-324, 2010.

Effect of Itch, Scratching and Mental Stress on Autonomic Nervous System Function in Atopic Dermatitis.
 Tran B.W., Papoiu A.D.P., Russoniello C.V., Wang H., Patel T.S.; Chan Y.-H. and Yosipovitch G.
Acta Derm Venereol, Jun; 90(4), 354-361, 2010.

5-Aminolevulinic Acid-based Photodynamic Intense Pulsed Light Therapy Shows Better Effects in the Treatment of Skin Photoaging in Asian Skin.
 Yang G., Xiang L.H. and Gold M.H.
J Clin Aesthet Dermatol, Mar; 3(3): 40-43, 2010.

A pramoxine-based anti-itch lotion is more effective than a control lotion for the treatment of uremic pruritus in adult hemodialysis patients.
 Young T.A., Patel T.S., Camacho F., Clark A., Freedman B.I., Kaur M., Fountain J., Williams L.L., Yosipovitch G. and Fleischer A.B. Jr.
J Dermatolog Treat, 20(2): 76-81, 2009.

Hydrating effects of moisturizer active compounds incorporated into hydrogels: in vivo assessment and comparison between devices
 Silva V.R.L., Schulman M.A., Ferelli C., Gimenes J.M., Ruas G.W., Baby A.R., Velasco M.V.R., Taqueda M.E. and Kaneko T.M.
J Cosmet Dermatol, published online 17 Feb 2009.

Noninvasive transcutaneous sampling of glucose by electroporation
 Murthy S.S., Kiran V.S.R., Mathur S.K. and Murthy S.N.
J Diabetes Sci Technol, 2(3):250-254, 2008.

The use of an extended ventilation tube as a countermeasure for EVA-associated upper extremity medical issues
 Jones J.A., Hoffman R.B., Buckland D.A., Harvey C.M., Bowen C.K., Huday C.E. Strauss S., Novak J. and Gernhardt M.L.
Acta Astronaut, 63:763-768, 2008.

Measurement of hydration in the stratum corneum with the MoistureMeter and comparison with the Corneometer
 Alanen E., Nuutinen J., Nicklén K., Lahtinen T. and Mönkkönen J.
Skin Res Technol, 10: 32-37, 2004.

Dielectric properties of the skin
 Lahtinen, T., Nuutinen, J., Alanen, E.
 In: Radio Frequency Radiation Dosimetry, Editors: B. J. Klauenberg and D. Miklavcic. Kluwer Academic Publishers, the Netherlands, 2000.

Penetration of electromagnetic fields of open-ended coaxial probe between 1 MHz and 1GHz in dielectric skin measurements
 Alanen, E., Lahtinen T. and Nuutinen J.
Phys Med Biol, 44(7): N169-176, 1999.

Variational formulation of open-ended coaxial line in contact with layered biological medium
 Alanen, E. Lahtinen, T. and Nuutinen, J.
IEEE Trans Biomed Eng, 45 (10) 1241-1248, 1998.

Dielectric properties of the skin
 Lahtinen, T., Nuutinen, J. and Alanen, E.
Phys Med Biol 42: 1471-1472, 1997.

Abstracts

Mechanism of injury and countermeasures for EVA associated upper extremity medical issues: Extended Vent Tube Study
Jones J., Hoffman R., Harvey G., Bowen C.K., Hudy C.E., Tuxhorn J., Gernhardt M. and Scheuring R.
Prenestation at 15th International Academy of Astronautics' "Humans in space" symposium in Beijing, 20-24 May 2007.

The moisturization ability of a sodium sulfacetamide (10%)/sulfur (5%) emollient foam
Trumbore M
J Am Acad Dermatol, February: AB21, 2007.

Water content at skin surface and in whole skin during irritant skin reaction
Miettinen M., Nuutilainen J., Lahtinen M.-R. and Lahtinen T.
13th Congress of European Academy of Dermatology and Venereology Meeting, 17-21 November, 2004.

Requirements for the conductivity measurements of the MoistureMeter
Alanen, E., Nuutilainen, J. and Nicklén, K.
Stratum Corneum IV. 17-19 June 2004: p. 48.

Susceptibility to SLS irritation: Effect of a five-day pre-use of two skin care oils studied comparing various techniques and instruments
Rantanen, T., Snellman, E. and Kostiainen M.
First World Congress on Work-Related and Environmental Allergy. 9-12 July 2003: p. 95.

Pressure-controlled measurement of skin surface hydration with MoistureMeter
Nuutilainen, J., Alanen, E. and Nicklén, K.
Skin Res Technol, Vol 9 (2): 202, 2003.